



# Soy Protein: A High Quality, Complete Protein

**S**oy protein is a high quality, complete plant protein that is comparable to high quality animal protein (e.g., beef, milk, eggs). As a primary source of protein, soy can help children and adults adequately meet their protein needs. Soy protein is unique in that it is the only complete source of vegetable protein widely


available. Protein could be considered the most important macronutrient for humans<sup>1,2</sup>. The roles of protein in the human body are numerous. Among other things, proteins function as enzymes and hormones, as transporters and receptors, and

as part of the structural integrity of most organs in the body<sup>2</sup>. Amino acids serve as precursors for nucleic acids, hormones, vitamins and other important molecules<sup>3</sup>. With so many important roles in all aspects of bodily functioning, it is essential that humans consume enough high-quality protein each day to meet the body's demands.


## Comparability of Different Proteins

The PDCAAS values for Solae™ isolated soy protein and other common proteins are presented below. These data show that soy protein achieves the maximum PDCAAS value of 1.00 and is deemed a nutritionally complete protein.

### PDCAAS Of Select Proteins



Sources: Solae In-house data, PDCAAS Value for Solae™ SUPRO® Brand Isolated Soy Protein; FAO/WHO (1991), Protein Quality Evaluation; FAO Food and Nutrition Paper 51, Rome, Italy.



**PDCAAS methodology takes into consideration three important parameters:**

- 1) the food protein's essential amino acid profile,
- 2) corrected for its digestibility
- 3) its ability to supply the FAO/WHO's amino acid, and requirements for 2-5 year olds<sup>4</sup>.

For humans, amino acid requirement values have been used to develop reference amino acid patterns for the purpose of evaluating the quality of food proteins or their capacity to efficiently meet both the nitrogen and indispensable amino acid requirements of an individual. Protein quality is a measure of protein bioavailability, and its evaluation is a means of determining the capacity of food proteins and diets to satisfy the metabolic demands of amino acids and nitrogen.

The Food and Agriculture Organization (FAO) and the World Health Organization (WHO) Joint Expert Consultation's Protein Quality Evaluation Report<sup>4</sup> recommended the protein digestibility-corrected amino acid score (PDCAAS), which is a simple, scientific and rational procedure for assessing protein quality. The United States Institute of Medicine (IOM) in its 2005 discussion of Daily Reference Intakes (DRIs) reaffirmed the use of PDCAAS for evaluating the relative nutritional quality of different protein sources<sup>3</sup>. Additionally, the US Food and Drug Administration (US FDA) adopted the use of PDCAAS for the protein quality evaluation of foods in 1993.

The PDCAAS methodology takes into consideration three important parameters: 1) the food protein's essential amino acid profile, 2) corrected for its digestibility and 3) its ability to supply the FAO/WHO's amino acid requirements for 2-5 year olds<sup>4</sup>. Although there are relatively small differences in amino acid requirements for age-sex groups beyond infancy, the 2-5 year old requirement is used because it is the most demanding pattern of any age group excluding that of infant requirements. For a given protein, the essential amino acid that is present in the relatively lowest quantity when compared to the standard is used for calculation of the final PDCAAS value. The highest possible score using the PDCAAS method is 1.00<sup>4</sup>.

The digestibility of protein is defined as the proportion of ingested protein that is absorbed. True digestibility of protein corrects for endogenous fecal nitrogen (measured as nitrogen loss that occurs on a protein-free diet) and is expressed as a true digestibility percent. The true digestibility of Solae isolated soy proteins is 97% (internal data), and comparable to other high quality proteins such as milk, meat and eggs that range between 94-97%<sup>5</sup>. Other plant proteins are of lower digestibility.

The use of Protein Efficiency Ratio (PER) to assess protein quality, although still used by some, measures the ability of a protein to support growth in young growing rats. It over-estimates the value of some animal proteins and underestimates the value of some vegetable proteins for human growth<sup>4,6</sup>. On the other hand, PDCAAS considers the full contribution of vegetable proteins to the diet. Additionally, vegetable proteins such as high quality soy protein have economic and sustainability advantages when compared to animal proteins.



**Table 1: Suggested Indispensable Amino Acid Requirements Published by the WHO/FAO<sup>5</sup> and the U.S. IOM<sup>3</sup> Expressed as Milligrams per Gram Protein**

Amino Acid	WHO/FAO Suggested Requirements			U.S. IOM	
	1 - 2 yrs	3 - 10 yrs	Adults	1 - 3 yrs	Adults
Histidine	18	16	15	18	17
Isoleucine	31	31	30	25	23
Leucine	63	61	59	55	52
Lysine	52	48	45	51	47
Methionine + Cysteine	26	24	22	25	23
Phenylalanine + Tyrosine	46	41	38	47	41
Threonine	27	25	23	27	24
Tryptophan	7.4	6.6	6.0	7	6
Valine	42	40	39	32	29

## Meeting Human Protein Requirements

Soy protein meets the essential amino acid requirements of children and adults. National and international organizations<sup>3-5,7</sup> and individual researchers<sup>8</sup> have published protein and amino acid requirement recommendations. These recommendations are used to assess whether a protein source is nutritionally complete based on its amino acid composition. The most recent estimations of the indispensable amino acid requirements for various age groups were published by the WHO/FAO<sup>5</sup>. Values for children and adults were published by the U.S.

Soy is a nutritionally complete protein with PDCAAS=1.00 and comparable to milk, eggs and meat.

IOM<sup>3</sup> as part of the DRIs for protein and amino acids. These values are presented above. Isolated soy proteins commercially available from Solae meet the published estimated indispensable

amino acid requirements for children and adults. Selected age groups are presented in Table 1.

Isolated soy protein has been shown to maintain nitrogen balance when fed as the sole protein source at minimum recommended intake levels. The nutritional characteristics of soy protein in human nutrition have been reviewed<sup>9-11</sup>. Soy protein was once widely regarded as nutritionally inferior to animal protein. The sulfur-containing amino acid methionine is the first limiting amino acid in soy protein, and animal bioassays such as the rat-based PER suggest that the biological value of soy protein is only 62-92% compared to that of casein. However, animal assays underestimate biological value because rats have a higher methionine requirement than humans<sup>9</sup> so these techniques cannot be used to accurately estimate the quality of soy protein for humans. Unequivocal data generated during the past 25 years have shown that soy is a nutritionally complete protein based on its indispensable amino acid spectrum.

Numerous nitrogen balance studies<sup>12-18</sup> have demonstrated that unmodified soy protein, when fed in nutritionally appropriate amounts to human subjects, is comparable in quality to that of beef, milk and other animal proteins. These nitrogen balance studies provide clear, consistent evidence that isolated soy protein (including Solae™ SUPRO® brand isolated soy protein) is a complete protein that can meet protein requirements. These data further provide direct, definitive evidence that isolated soy protein is nutritionally comparable to high quality animal proteins when fed in nutritionally appropriate amounts.





## REFERENCES

1. Reeds P, Schaafsma G, Tome D, Young V (2000). Criteria and significance of dietary protein sources in humans; summary of the workshop with recommendations. *J Nutr*, 130:1874S-6S.
2. Otten JJ, Hellwig JP, Meyers LD (2006). *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: The National Academies Press.
3. Institute of Medicine (2005). *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids*, Washington, DC: The National Academies Press.
4. FAO/WHO (1991). *Protein Quality Evaluation*; FAO Food and Nutrition Paper 51, Rome, Italy.
5. WHO/FAO/UNU (2007). *Protein and Amino Acid Requirements in Human Nutrition*. Report of a Joint WHO/FAO/UNU Expert Consultation, Geneva, Switzerland: World Health Organization.
6. Henley EC, Kuster JM (1994). Protein quality evaluation by protein digestibility-corrected amino acid scoring. *Food Technology*, 48:74-7.
7. FAO/WHO/UNU (1985). *Energy and Protein Requirements*. Report of a Joint FAO/WHO/UNU Expert Consultation, Geneva, Switzerland: World Health Organization.
8. Young V & Pellett P (1990). Current concepts concerning indispensable amino acid needs in adults and their implications for international nutrition planning. *Food Nutr Bull*, 12: 289-300.
9. Erdman JW Jr & Fordyce EJ (1989). Soy products and the human diet. *Am J Clin Nutr*, 49: 725-37.
10. Friedman M & Brandon DL (2001). Nutritional and health benefits of soy proteins. *J Agric Food Chem*, 49: 1069-86.
11. Jeejeebhoy KN (2000). Vegetable proteins: are they nutritionally equivalent to animal protein. *Eur J Gastroenterol Hepatol*, 12: 1-2.
12. Zezulka AY & Calloway DH (1976). Nitrogen retention in men fed isolated soybean protein supplemented with L-methionine, D-methionine, N-acetyl-L-methionine, or inorganic sulfate. *J Nutr*, 106: 1286-91.
13. Istfan N, Murray E, Janghorbani M & Young, VR (1983). An evaluation of the nutritional value of a soy protein concentrate in young adult men using the short-term N-balance method. *J Nutr*, 113: 2516-23.
14. Scrimshaw NS, Wayler AH, Murray E, Steinke FH, Rand WM & Young VR (1983). Nitrogen balance response in young men given one of two isolated soy proteins or milk proteins. *J Nutr*, 113: 2492-7.
15. Wayler A, Queiroz E, Scrimshaw NS, Steinke FH, Rand WM & Young VR (1983). Nitrogen balance studies in young men to assess the protein quality of an isolated soy protein in relation to meat proteins. *J Nutr*, 113: 2485-91.
16. Young VR, Wayler A, Garza C, Steinke FH, Murray E, Rand WM & Scrimshaw NS (1984b). A long-term metabolic balance study in young men to assess the nutritional quality of an isolated soy protein and beef proteins. *Am J Clin Nutr*, 39: 8-15.
17. Young VR, Puig M, Queiroz E, Scrimshaw NS & Rand WM (1984a). Evaluation of the protein quality of an isolated soy protein in young men: relative nitrogen requirements and effect of methionine supplementation. *Am J Clin Nutr*, 39: 16-24.
18. Beer WH, Murray E, Oh SH, Pedersen HE, Wolfe RR & Young VR (1989). A long-term metabolic study to assess the nutritional value of and immunological tolerance to two soy-protein concentrates in adult humans. *Am J Clin Nutr*, 50: 997-1007.

*The information contained herein is, to the best of our knowledge, correct. The data outlined and the statements made are intended only as a source of information. Also, we may suggest technical solutions for incorporating this ingredient into products, however, it is the user's responsibility to comply with appropriate government standards and requirements. No warranties, expressed or implied, are made. On the basis of this information, it is suggested that you evaluate the product on a laboratory scale prior to use in a finished product. The information contained herein should not be construed as permission for violation of patent rights.*

**Solae**<sup>TM</sup>

4300 Duncan Avenue | St. Louis, Missouri 63110  
1.800.325.7108 | [www.solae.com](http://www.solae.com)

© Solae, LLC 2009. All rights reserved.  
All products denoted with <sup>TM</sup> or ® are trademarks or registered trademarks of Solae, LLC and/or its affiliates.

